

REPORT DOCUMENTATION PAGE*Form Approved*
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY)

15-06-2001

2. REPORT TYPE

UU

3. DATES COVERED (From - To)**4. TITLE AND SUBTITLE**

See the abstract below

5a. CONTRACT NUMBER**5b. GRANT NUMBER****5c. PROGRAM ELEMENT NUMBER****6. AUTHOR(S)**

See the abstract below

5d. PROJECT NUMBER**5e. TASK NUMBER****5f. WORK UNIT NUMBER****7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)****8. PERFORMING ORGANIZATION REPORT NUMBER****9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)****10. SPONSOR/MONITOR'S ACRONYM(S)****11. SPONSOR/MONITOR'S REPORT NUMBER(S)****12. DISTRIBUTION / AVAILABILITY STATEMENT****DISTRIBUTION STATEMENT A**
Approved for Public Release
Distribution Unlimited

20010705061

13. SUPPLEMENTARY NOTES**14. ABSTRACT**

Phone.txt



DTICPUB0615.txt

US NAVAL OBSERVATORY COLLECTION

15. SUBJECT TERMS**16. SECURITY CLASSIFICATION OF:**

UU

17. LIMITATION OF ABSTRACT**18. NUMBER OF PAGES****19a. NAME OF RESPONSIBLE PERSON**
See the abstract above

a. REPORT

b. ABSTRACT

c. THIS PAGE

19b. TELEPHONE NUMBER (include area code)
See the phone list above

Beish, J. D. (1999). "Windows International Mars Patrol Astronomical Calculator WIMP," The Digital Lens: Newsletter of the Association of Lunar and Planetary Observers Computing Section,
<http://www.m2c3.com/alpocs/tcl1999/wimpinfo10031999/WIMP.html>

Beish, J. D., (2000). "Discrete Topographic and Orographic Clouds of Mars," Association of Lunar and Planetary Observers Internet Web Page: The Mars Section,
<http://www.lpl.arizona.edu/~rhill/alpo/marstuff/discrete.htm>

Beish, J. D. (2000). "A German Equatorial Mount for the Planetary Telescope," The Digital Lens: Newsletter of the Association of Lunar and Planetary Observers Computing Section,
<http://www.m2c3.com/alpocs/tcl2000/mountmath05222000/mount.htm>

Beish, J. D. (2000). "Martian Volcanoes on HST Images " Association of Lunar and Planetary Observers Internet Web Page: The Mars Section,
<http://www.lpl.arizona.edu/~rhill/alpo/marstuff/VOLCANO.htm>

Beish, J. D. (2000). "Measuring Celestial Dimensions with Micrometers," The Digital Lens: Newsletter of the Association of Lunar and Planetary Observers Computing Section,
http://www.m2c3.com/alpocs/tcl2000/celestialdimensions01232000/celestial_dimensions.htm

Beish, J. D. (2000). "Nuts and Bolts of Computing the Ephemeris - Part Four," The Digital Lens: Newsletter of the Association of Lunar and Planetary Observers Computing Section,
<http://www.m2c3.com/alpocs/tcl1999/nutsbolts08301999/TDL-4.html>

Beish, J.D. (2000). "Practical Calculations for the Newtonian Secondary Mirror," The Digital Lens: Newsletter of the Association of Lunar and Planetary Observers Computing Section,
<http://www.m2c3.com/alpocs/tcl2000/telescope/math06202000/field1.htm>

Beish, J. D. (2000). "Some of the 'Astronomical Seeing' Scales," The Digital Lens: Newsletter of the Association of Lunar and Planetary

Discrete Topographic and Orographic Clouds of Mars

DISTRIBUTION STATEMENT A

By: Jeff Beish

Approved for Public Release Association of Lunar and Planetary Observers (A.L.P.O.)

Distribution Unlimited

INTRODUCTION

The Martian atmosphere is very dynamic, exhibiting several types of clouds and hazes easily detected with the aid of color filters and modest-sized telescopes. White water clouds, local yellowish dust clouds, global dust storms, bluish limb hazes, and bright surface *ice-fogs* and frosts have been studied with increased interest over the past three decades. Observations of these meteorological features indicate their behavior and occurrence are most often coupled with the seasonal sublimation and condensation of polar-cap material.

Although the Martian sky appears quite clear, transient bright streaks and oval patches are often observed in its atmosphere that exhibits similar characteristics to clouds and hazes seen in the Earth's atmosphere. Fine dust particles, CO₂ and/or H₂O hazes and ice crystals are regularly observed in Mars' atmosphere using moderate telescopes. From the data gathered by the Mariner and Viking Land-er/Orbiter spacecraft, we know that H₂O and CO₂ condensates do exist in the atmosphere of Mars. We are comfortable with the idea that -- what we observe, as bright patches on Mars are in fact atmospheric clouds, fogs, and/or hazes.

In the Earth's atmosphere clouds often form on the windward side of mountains and are classified as Orographic Clouds. Orographic clouds are clouds that develop in response to the forced lifting of air by the earth's topography, such as mountains, and fall into the following general categories:

Conjoined orographic clouds - covers entire mountain, forms on windward side.

Orographic Cumulus fractus - usually covering mountain range, large area of clouds covering both windward and lee sides of mountains.

Orographic Stratocumulus and Altocumulus (wave clouds) - form on the lee side of a mountain.

Pileus (Latin for "skullcap") is a smooth cloud found attached to either a mountaintop or growing cumulus tower.

HOW DISCRETE CLOUDS BEHAVE

Discrete Topographic and Orographic Clouds tend to rotate with the diurnal rotation of the planet. These type phenomena are identified by their location near historically cloudy regions and high mountainous areas. When comparing the historical trends [*Beish et al*, 1986], [*Beish et al*, 1987], [*Beish and Parker*, 1990], and [*Beish* 1999], it is evident that discrete clouds are more numerous during the northern summer and appears to follow a similar seasonal trend line to both Evening and Morning clouds. As the northern summer begins discrete clouds continue to increase in number and frequency after limb clouds begin decreasing and peak during northern mid-summer. Their number and frequency slowly drops off in late northern summer throughout the remainder of the Martian year.

These discrete white clouds are especially prominent in mountainous regions and volcanoes after rapid

20010705 061

AOI 01-16 -1797

thawing of the North Polar Cap (NPC) begins. By early afternoon, they appear as bright round or oval-shaped individual clouds and at times producing a "Domino Effect" on the Martian disk. Cloud formations can be seen without filters; however, they tend to brighten in blue light. They usually continue to expand and brighten as the planet rotates toward the sunset limb and will brighten violet light, indicating that they also expand upwards by convection. Some of these clouds coalesce with other clouds to form large and bright limb clouds such as the well-known "W-clouds" observed in Tharsis-Amazonis region. A particularly bright section of the W-clouds can be seen over Nix Olympica or Olympus Mons (113°W, 18°N).

Other areas for cloud formations are Nix Tanaica (55°W, 52°N), Nix Cydonia (218°W, 00°N, Arsia Mons (°121W, 19°S), Apollinaris Petera (186°W, 08°S), Ascraeus Mons (104°W, 11°N), Pavonis Mons (112°W, 0°N), and Arisa Mons (120°W, 9°S), and Elysium Mons (212°W, 25°N), and between Tharsis Tholus and Valles Marineris (80°-100°, 04°N). These cloud formations are usually groups of orographic clouds.

Orographic Clouds. On Earth, large dense clouds often form in mountainous regions and hover over the tops of mountains. Air that flows over a mountain forms airwaves on the leeward side of the mountain. In the back of the wave where the air rises, cools and forms smooth elongated clouds. When moist air reaches colder upper altitudes rapidly cooling causes condensation to occur and clouds form. These clouds will linger over mountainous regions for long periods and move around very little in spite of very strong winds. They are referred to as *orographic clouds*. Orographic clouds on Earth are often called almond clouds or lenticular clouds [Miller and Thompson, 1970].

Martian orographic clouds exhibit similar characteristics. Orographic clouds form when the Martian atmosphere rises while flowing up the slopes and over mountains to begin the condensation process. Spacecraft images suggested clouds were lee-wave clouds following the large volcanoes and where found in many regions of the Martian surface. When the gaseous constituents of Mars' atmosphere flow over a mountain and form waves on the leeward side of the mountain, it cools and forms a cloud. Unlike clouds in earth's thicker atmosphere that form into water droplets or a mist, Martian clouds condense directly from vapor to ice particles. The atmospheric pressure on Mars is just too low for liquid water to form.

Topographic Clouds. Dense bluish-white clouds of limited extent populate many areas of Mars during spring and summer after persisting for days in the same general areas. Topographic clouds appear in or near large, deep craters and over great plains and valleys. Three areas; Aram (12°W, 05°S), Edom (345°W, 04°S), and Ophir (65°W, 10°S), are particularly fertile sites for white cloud formation. Another region has been observed to be cloudy during each apparition is the Isidis Regio-Libya-Crocea (275°W, 20°N - 293°W, 00°N) near Syrtis Major. Yellowish clouds, blue-white clouds, and low hazes seem to gravitate into this large depression that lies north of the Tyrrenum-Hesperia (240°W, 20°S) block and just east of the up-slopes of the Syrtis Major Planum (290°W, 10°N). Wave clouds form on the sheltered side of large obstacles, such as craters, and have very distinct ridges.

Topographic clouds show growth and displacement within a region, especially when near the sunset limb. Often they are called "localized clouds" or "regional discrete clouds" and appear as white patches. They will brighten when observed in blue or blue-green light indicating they are possibly blue or blue-white clouds. Their color can vary; they can become prominent in green or can brighten in yellow light. This led C.F. Capen and the authors to believe they are a combination of water vapor and dust. Topographic clouds are usually long and/or irregular shaped blue or white formations found in bright desert regions. However, they are often observed over portions of dark 'maria' as well. Clouds of this type follow the rotation of the planet and are often observed for several Martian days in the same general

areas. These clouds sometimes form large wedge shapes and may be found in regions where the low plains meet higher elevations, such as the region southeast of the Tharsis bulge.

DISCUSSION

The International Mars Patrol (I.M.P.) has initiated an observing program for intensive investigation into these phenomena and will appeal to all planetary observers using CCD technology to assist us in this important study.

The Hubble Space Telescope (HST) has revealed that these clouds follow a similar trend line or forecast predicted in recent A.L.P.O. meteorological studies of Mars.

Topographic and Orographic Clouds are best detected visually through a light blue (W38A or 80A) and deep blue (W47 and W47B) Wratten filters and may be photographed or imaged in blue or ultraviolet light. They are sometimes observed quite easily in integrated light (non-filters) as well.

New technologies, such as CCD cameras, sophisticated computer hardware and software, and large-aperture planetary telescopes have given rise to a virtual explosion in advanced techniques of studying our Solar System. Never before have we been able to readily detect the delicate wispy Martian Topographic and Orographic Clouds so well as we do now with CCD imaging.

While it is not the purpose to discuss the physics of these clouds we never the less draw certain conclusions from observational evidence. Questions remain that may be answered with high-tech observing techniques by amateur astronomers. There is no reason we cannot at least serve as long termed clouds reporters between scant times of HST observing periods.

REFERENCES

Beish, J.D., Parker, D.C., and Capen, C.F., "Meteorology of Mars - Part I," *Journal of the Association of Lunar and Planetary Observers (J.A.L.P.O.)*, Vol.31, Nos. 11-12, November 1986.

Beish, J.D., Parker, D.C., and Capen, C.F., "Meteorology of Mars - Part II," *J.A.L.P.O.*, Vol.32, Nos. 1-2, March 1987.

Beish, J.D. and Parker, D.C., "Meteorology of Mars - Part III," *J.A.L.P.O.*, Vol. 32, Nos. 5-6, October 1987.

Beish, J.D., and D.C. Parker, "Meteorological Survey of Mars, 1968-1985," *Journal of Geophysical Research*, Vol. 95, B9, 14657-14675, August 20, 1990.

Beish, J.D., "Meteorological Survey of Mars For Opposition Years 1965 - 1995," *The ALPO Internet Web Page: The Digital Lens*, December 1999.

<http://www.m2c3.com/alpocs/tdl1999/meteorological110199/MOM.html>

Miller, Albert, and Jack C., **Thompson, Elements of Meteorology**, Merrill Publishing, Columbus, Ohio 78-98476, 1970.